

## Standard Recommendation Language

---

**A**—The USPSTF strongly recommends that clinicians routinely provide [the service] to eligible patients. (The USPSTF found good evidence that [the service] improves important health outcomes and concludes that benefits substantially outweigh harms.)

**B**—The USPSTF recommends that clinicians routinely provide [the service] to eligible patients. (The USPSTF found at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms.)

**C**—The USPSTF makes no recommendation for or against routine provision of [the service]. (The USPSTF found at least fair evidence that [the service] can improve health outcomes but concludes that the balance of the benefits and harms is too close to justify a general recommendation.)

**D**—The USPSTF recommends against routinely providing [the service] to asymptomatic patients. (The USPSTF found at least fair evidence that [the service] is ineffective or that harms outweigh benefits.)

**I**—The USPSTF concludes that the evidence is insufficient to recommend for or against routinely providing [the service]. (Evidence that [the service] is effective is lacking, of poor quality, or conflicting and the balance of benefits and harms cannot be determined.)

---

**Note:** All statements specify the population for which the recommendation is intended and are followed by a rationale statement providing information about the overall grade of evidence and the net benefit from implementing the service.

**Source:** [Third U.S. Preventive Services Task Force \(USPSTF\)](#).

Source: <http://www.ahrq.gov/clinic/uspstf/standard.htm>